

EndNote Web

Search Key Word: Pulsed field gel electrophoresis or PFGE and Neisseria gonorrhoeae (in Abstract)

Search date: August 13, 2008

Export in the bibliographic style of J Clin Microbiol

1. **Azariah, S., and N. Perkins.** 2007. Risk factors and characteristics of patients with gonorrhoea presenting to Auckland Sexual Health Service, New Zealand. *N Z Med J* **120**:U2491.
2. **Billings, S., D. Fuller, A. LeMonte, T. Davis, and A. Hartstein.** 1997. Characterization of DFA-negative, probe-positive Neisseria gonorrhoeae by pulsed field gel electrophoresis. *Diagn Microbiol Infect Dis* **29**:281-3.
3. **Blackmore, T., G. Hererra, S. Shi, P. Bridgewater, L. Wheeler, and J. Byrne.** 2005. Characterization of prolyl iminopeptidase-deficient Neisseria gonorrhoeae. *J Clin Microbiol* **43**:4189-90.
4. **Chang, N., and L. Chui.** 1998. A standardized protocol for the rapid preparation of bacterial DNA for pulsed-field gel electrophoresis. *Diagn Microbiol Infect Dis* **31**:275-9.
5. **Cooke, S., H. de la Paz, C. La Poh, C. Ison, and J. Heckels.** 1997. Variation within serovars of Neisseria gonorrhoeae detected by structural analysis of outer-membrane protein PIB and by pulsed-field gel electrophoresis. *Microbiology* **143 (Pt 4)**:1415-22.
6. **Corkill, J., A. Komolafe, T. Neal, A. Mortimore, A. Alawattegama, and C. Hart.** 2003. Molecular epidemiology of endemic ciprofloxacin-resistant Neisseria gonorrhoeae in Liverpool. *Int J STD AIDS* **14**:379-85.
7. **DeMattia, A., J. Kornblum, J. Hoffman-Rosenfeld, D. Trees, A. Tumpey, and D. Laraque.** 2006. The use of combination subtyping in the forensic evaluation of a three-year-old girl with gonorrhea. *Pediatr Infect Dis J* **25**:461-3.
8. **Dempsey, J., and J. Cannon.** 1994. Locations of genetic markers on the physical map of the chromosome of Neisseria gonorrhoeae FA1090. *J Bacteriol* **176**:2055-60.
9. **Dempsey, J., A. Wallace, and J. Cannon.** 1995. The physical map of the chromosome of a serogroup A strain of Neisseria meningitidis shows complex rearrangements relative to the chromosomes of the two mapped strains of the closely related species N. gonorrhoeae. *J Bacteriol* **177**:6390-400.
10. **Fjeldsøe-Nielsen, H., M. Unemo, H. Fredlund, S. Hjorth, L. Berthelsen, H. Palmer, and A. Friis-Møller.** 2005. Phenotypic and genotypic characterization of prolyliminopeptidase-negative Neisseria gonorrhoeae isolates in Denmark. *Eur J Clin Microbiol Infect Dis* **24**:280-3.
11. **Garvin, L., M. Bash, C. Keys, D. Warner, S. Ram, W. Shafer, and A. Jerse.** 2008. Phenotypic and genotypic analyses of Neisseria gonorrhoeae isolates that express frequently recovered PorB PIA variable region types suggest that certain P1a porin sequences confer a selective advantage for urogenital tract infection. *Infect Immun* **76**:3700-9.
12. **Gordon, S., C. Carlyn, L. Doyle, C. Knapp, D. Longworth, G. Hall, and J. Washington.** 1996. The emergence of Neisseria gonorrhoeae with decreased susceptibility to ciprofloxacin in Cleveland, Ohio: epidemiology and risk factors. *Ann*

- Intern Med **125**:465-70.
13. **Gunn, J., A. Piekarowicz, R. Chien, and D. Stein.** 1992. Cloning and linkage analysis of *Neisseria gonorrhoeae* DNA methyltransferases. *J Bacteriol* **174**:5654-60.
 14. **Gäher, M., K. Einsiedler, T. Crass, and W. Bautsch.** 1996. A physical and genetic map of *Neisseria meningitidis* B1940. *Mol Microbiol* **19**:249-59.
 15. **Hamrick, T., J. Dempsey, M. Cohen, and J. Cannon.** 2001. Antigenic variation of gonococcal pilin expression in vivo: analysis of the strain FA1090 pilin repertoire and identification of the pilS gene copies recombining with pilE during experimental human infection. *Microbiology* **147**:839-49.
 16. **Harnett, N., S. Brown, G. Riley, R. Terro, C. Krishnan, M. Pauzé, and K. Yeung.** 1997. Analysis of *Neisseria gonorrhoeae* in Ontario, Canada, with decreased susceptibility to quinolones by pulsed-field gel electrophoresis, auxotyping, serotyping and plasmid content. *J Med Microbiol* **46**:383-90.
 17. **Harnett, N., S. Brown, R. Terro, C. Krishnan, M. Pauzé, and K. Yeung.** 1997. High-level tetracycline-resistant *Neisseria gonorrhoeae* in Ontario, Canada--investigation of a cluster of isolates, showing chromosomally mediated resistance to penicillin combined with plasmid-mediated resistance to tetracycline. *J Infect Dis* **176**:1269-76.
 18. **Hesse, S., N. Low, B. Molokwu, A. Uttley, A. Jephcott, A. Turner, and A. Pozniak.** 1995. Molecular typing as an epidemiological tool in the study of sexual networks. *Lancet* **346**:976-7.
 19. **Hsueh, P., S. Tseng, L. Teng, and S. Ho.** 2005. High prevalence of ciprofloxacin-resistant *Neisseria gonorrhoeae* in Northern Taiwan. *Clin Infect Dis* **40**:188-92.
 20. **Ito, M., T. Deguchi, K. Mizutani, M. Yasuda, S. Yokoi, S. Ito, Y. Takahashi, S. Ishihara, Y. Kawamura, and T. Ezaki.** 2005. Emergence and spread of *Neisseria gonorrhoeae* clinical isolates harboring mosaic-like structure of penicillin-binding protein 2 in Central Japan. *Antimicrob Agents Chemother* **49**:137-43.
 21. **Iyoda, T., T. Saika, A. Kanayama, M. Hasegawa, I. Kobayashi, Y. Onoe, M. Tanaka, and S. Naito.** 2003. [Bacteriological and epidemiological study on *Neisseria gonorrhoeae* isolated from the pharyngeal specimens of male and female patients with gonorrhoea]. *Kansenshogaku Zasshi* **77**:103-9.
 22. **Kilmarx, P., J. Knapp, M. Xia, M. St Louis, S. Neal, D. Sayers, L. Doyle, M. Roberts, and W. Whittington.** 1998. Intercity spread of gonococci with decreased susceptibility to fluoroquinolones: a unique focus in the United States. *J Infect Dis* **177**:677-82.
 23. **Lee, K., Y. Chong, L. Erdenechemeg, K. Soon Song, and K. Hun Shin.** 1998. Incidence, epidemiology and evolution of reduced susceptibility to ciprofloxacin in *Neisseria gonorrhoeae* in Korea. *Clin Microbiol Infect* **4**:627-633.
 24. **Lee, K., J. Shin, J. Lim, Y. Kim, D. Yong, H. Oh, and Y. Chong.** 2000. Emerging antimicrobial resistance, plasmid profile and pulsed-field gel electrophoresis pattern of the endonuclease-digested genomic DNA of *Neisseria gonorrhoeae*. *Yonsei Med J* **41**:381-6.
 25. **Li, H., and J. Dillon.** 1995. Utility of ribotyping, restriction endonuclease analysis and pulsed-field gel electrophoresis to discriminate between isolates of *Neisseria gonorrhoeae* of serovar IA-2 which require arginine, hypoxanthine or uracil for growth. *J Med Microbiol* **43**:208-15.
 26. **Lum, G., K. Freeman, N. Nguyen, E. Limnios, S. Tabrizi, I. Carter, I. Chambers, D.**

- Whiley, T. Sloots, S. Garland, and J. Tapsall.** 2005. A cluster of culture positive gonococcal infections but with false negative *cppB* gene based PCR. *Sex Transm Infect* **81**:400-2.
27. **Lundbäck, D., H. Fredlund, T. Berglund, B. Wretling, and M. Unemo.** 2006. Molecular epidemiology of *Neisseria gonorrhoeae*- identification of the first presumed Swedish transmission chain of an azithromycin-resistant strain. *APMIS* **114**:67-71.
28. **Mavroidi, A., L. Tzouveleki, P. Tassios, A. Flemetakis, M. Daniilidou, and E. Tzelepi.** 2000. Characterization of *Neisseria gonorrhoeae* strains with decreased susceptibility to fluoroquinolones isolated in Greece from 1996 to 1999. *J Clin Microbiol* **38**:3489-91.
29. **Ng, L., M. Carballo, and J. Dillon.** 1995. Differentiation of *Neisseria gonorrhoeae* isolates requiring proline, citrulline, and uracil by plasmid content, serotyping, and pulsed-field gel electrophoresis. *J Clin Microbiol* **33**:1039-41.
30. **Ng, L., and I. Martin.** 2005. The laboratory diagnosis of *Neisseria gonorrhoeae*. *Can J Infect Dis Med Microbiol* **16**:15-25.
31. **Ng, L., P. Sawatzky, I. Martin, and S. Booth.** 2002. Characterization of ciprofloxacin resistance in *Neisseria gonorrhoeae* isolates in Canada. *Sex Transm Dis* **29**:780-8.
32. **Poh, C., and Q. Lau.** 1993. Subtyping of *Neisseria gonorrhoeae* auxotype-serovar groups by pulsed-field gel electrophoresis. *J Med Microbiol* **38**:366-70.
33. **Poh, C., Q. Lau, and V. Chow.** 1995. Differentiation of *Neisseria gonorrhoeae* IB-3 and IB-7 serovars by direct sequencing of protein IB gene and pulsed-field gel electrophoresis. *J Med Microbiol* **43**:201-7.
34. **Poh, C., G. Loh, and J. Tapsall.** 1995. Resolution of clonal subgroups among *Neisseria gonorrhoeae* IB-2 and IB-6 serovars by pulsed-field gel electrophoresis. *Genitourin Med* **71**:145-9.
35. **Poh, C., V. Ramachandran, and J. Tapsall.** 1996. Genetic diversity of *Neisseria gonorrhoeae* IB-2 and IB-6 isolates revealed by whole-cell repetitive element sequence-based PCR. *J Clin Microbiol* **34**:292-5.
36. **Saika, T., T. Nishiyama, A. Kanayama, I. Kobayashi, H. Nakayama, M. Tanaka, and S. Naito.** 2001. Comparison of *Neisseria gonorrhoeae* isolates from the genital tract and pharynx of two gonorrhea patients. *J Infect Chemother* **7**:175-9.
37. **Sosa, J., S. Ramirez-Arcos, M. Ruben, H. Li, R. Llanes, A. Llop, and J. Dillon.** 2003. High percentages of resistance to tetracycline and penicillin and reduced susceptibility to azithromycin characterize the majority of strain types of *Neisseria gonorrhoeae* isolates in Cuba, 1995-1998. *Sex Transm Dis* **30**:443-8.
38. **Starnino, S., A. Neri, and P. Stefanelli.** 2008. Molecular analysis of tetracycline-resistant gonococci: rapid detection of resistant genotypes using a real-time PCR assay. *FEMS Microbiol Lett.*
39. **Su, X., N. Hutapea, J. Tapsall, and I. Lind.** 2003. Plasmid-mediated resistance of *Neisseria gonorrhoeae* strains isolated from female sex workers in North Sumatra, Indonesia, 1996. *Sex Transm Dis* **30**:178-82.
40. **Su, X., and I. Lind.** 2001. Molecular basis of high-level ciprofloxacin resistance in *Neisseria gonorrhoeae* strains isolated in Denmark from 1995 to 1998. *Antimicrob Agents Chemother* **45**:117-23.
41. **Tanaka, M., H. Nakayama, M. Haraoka, T. Nagafuji, T. Saika, and I. Kobayashi.** 1998. Analysis of quinolone resistance mechanisms in a sparfloxacin-resistant clinical

- isolate of *Neisseria gonorrhoeae*. *Sex Transm Dis* **25**:489-93.
42. **Tsunoe, H., M. Tanaka, O. Mochida, H. Nakayama, A. Kanayama, T. Saika, I. Kobayashi, and S. Naito.** 2001. Association between proline-requiring auxotype and fluoroquinolone resistance in *Neisseria gonorrhoeae* isolated in Japan. *J Infect Chemother* **7**:218-23.
 43. **Unemo, M., T. Berglund, P. OlcEn, and H. Fredlund.** 2002. Pulsed-field gel electrophoresis as an epidemiologic tool for *Neisseria gonorrhoeae*: identification of clusters within serovars. *Sex Transm Dis* **29**:25-31.
 44. **Unemo, M., P. Olcén, T. Berglund, J. Albert, and H. Fredlund.** 2002. Molecular epidemiology of *Neisseria gonorrhoeae*: sequence analysis of the *porB* gene confirms presence of two circulating strains. *J Clin Microbiol* **40**:3741-9.
 45. **Unemo, M., H. Palmer, T. Blackmore, G. Herrera, H. Fredlund, A. Limnios, N. Nguyen, and J. Tapsall.** 2007. Global transmission of prolyliminopeptidase-negative *Neisseria gonorrhoeae* strains: implications for changes in diagnostic strategies. *Sex Transm Infect* **83**:47-51.
 46. **Unemo, M., A. Sjöstrand, M. Akhras, B. Gharizadeh, E. Lindbäck, N. Pourmand, B. Wretlind, and H. Fredlund.** 2007. Molecular characterization of *Neisseria gonorrhoeae* identifies transmission and resistance of one ciprofloxacin-resistant strain. *APMIS* **115**:231-41.
 47. **Van Looveren, M., C. Ison, M. Ieven, P. Vandamme, I. Martin, K. Vermeulen, A. Renton, and H. Goossens.** 1999. Evaluation of the discriminatory power of typing methods for *Neisseria gonorrhoeae*. *J Clin Microbiol* **37**:2183-8.
 48. **Van Looveren, M., P. Vandamme, W. Wuyts, M. Ieven, and H. Goossens.** 2000. Organization of the ribosomal operon 165-235 gene spacer region in representatives of *Neisseria gonorrhoeae*. *Syst Appl Microbiol* **23**:9-14.
 49. **Xia, M., M. Roberts, W. Whittington, K. Holmes, J. Knapp, J. Dillon, and T. Wi.** 1996. *Neisseria gonorrhoeae* with decreased susceptibility to ciprofloxacin: pulsed-field gel electrophoresis typing of strains from North America, Hawaii, and the Philippines. *Antimicrob Agents Chemother* **40**:2439-40.
 50. **Xia, M., W. Whittington, K. Holmes, F. Plummer, and M. Roberts.** 1995. Pulsed-field gel electrophoresis for genomic analysis of *Neisseria gonorrhoeae*. *J Infect Dis* **171**:455-8.
 51. **Xia, M., W. Whittington, K. Holmes, and M. Roberts.** 1997. Genomic homogeneity of the AHU/IA-1,2 phenotype of *Neisseria gonorrhoeae* during its disappearance from an urban population. *Sex Transm Dis* **24**:561-6.
 52. **Xia, M., W. Whittington, W. Shafer, and K. Holmes.** 2000. Gonorrhea among men who have sex with men: outbreak caused by a single genotype of erythromycin-resistant *Neisseria gonorrhoeae* with a single-base pair deletion in the *mtrR* promoter region. *J Infect Dis* **181**:2080-2.
 53. **Yagupsky, P., A. Schahar, N. Peled, N. Porat, R. Trefler, M. Dan, Y. Keness, and C. Block.** 2002. Increasing incidence of gonorrhea in Israel associated with countrywide dissemination of a ciprofloxacin-resistant strain. *Eur J Clin Microbiol Infect Dis* **21**:368-72.
 54. **Yokoi, S., T. Deguchi, T. Ozawa, M. Yasuda, S. Ito, Y. Kubota, M. Tamaki, and S. Maeda.** 2007. Threat to cefixime treatment for gonorrhea. *Emerg Infect Dis* **13**:1275-7.
 55. **Yong, D., T. Kim, J. Choi, J. Yum, K. Lee, Y. Chong, H. Oh, T. Shultz, and J.**

- Tapsall.** 2004. Epidemiological characteristics and molecular basis of fluoroquinolone-resistant *Neisseria gonorrhoeae* strains isolated in Korea and nearby countries. *J Antimicrob Chemother* **54**:451-5.
56. **Yoo, J., C. Yoo, Y. Cho, H. Park, H. Oh, and W. Seong.** 2004. Antimicrobial resistance patterns (1999-2002) and characterization of ciprofloxacin-resistant *Neisseria gonorrhoeae* in Korea. *Sex Transm Dis* **31**:305-10.